

**Table 3J-14:
Acreage of Impacts to Sensitive Species Potentially Occurring in the
Project Area**

Species	Habitat Impacts (acres)		Determination of Effect ^a
	Alt. 2	Alt. 3	
BOREAL OWL (<i>Aegolius funereus</i>) (nesting/foraging)	27.6	23.8	MII
FISHER (<i>Martes pennanti</i>)	27.6	22.1	MII
FLAMMULATED OWL (<i>Otus flammeolus</i>) (nesting/foraging)	27.6	23.8	MII
GREATER SAGE-GROUSE (<i>Centrocercus urophasianus</i>)	1.2	1.2	No Impact
NORTHERN GOSHAWK (<i>Accipiter gentilis</i>) (nesting/foraging)	28.8	23.3	MII
NORTHERN THREE-TOED WOODPECKER (<i>Picoides tridactylus</i>) (nesting/foraging)	27.6	23.8	MII
PEREGRINE FALCON (<i>Falco peregrinus</i>)	1.2	1.2	MII
SPOTTED BAT (<i>Euderma maculatum</i>) (foraging/roosting)	1.2	1.2	MII
SPOTTED FROG (<i>Rana pretiosa</i>)	0	0	No Impact
TOWNSEND'S BIG-EARED BAT (<i>Corynorhinus townsendii</i>) (foraging/roosting)	1.2	1.2	MII
WOLVERINE (<i>Gulo gulo</i>) (denning/foraging/travel)	40.1	38.4	MII
WOOD RIVER SCULPIN (<i>Cottus leiopomus</i>)	0	0	MII
BLACK-BACKED WOODPECKER (<i>Picoides arcticus</i>)	38.9	15.7	MII
LEWIS' WOODPECKER (<i>Melanerpes lewis</i>)	28	7.3	MII
FRINGED MYOTIS (<i>Myotis thysanodes</i>)	1.2	1.2	MII

a - The determination of effect considers all impacts to the species and its habitat resulting from Ski Area Development and the 2005 VMP.

Note: MII = May impact individuals but would not likely contribute to a trend towards federal listing

